

LIVERPOOL SCHOOL OF TROPICAL MEDICINE



CLINICAL DIAGNOSTIC PARASITOLOGY LABORATORY

LABORATORY USER HANDBOOK 2020

Clinical Diagnostic Parasitology Laboratory,
Liverpool School of Tropical Medicine,
Pembroke Place,
Liverpool,
L3 5QA.

LSTM Diagnostic Laboratory,
DX 6966301
Liverpool 92L

<http://www.lstmed.ac.uk/CDPL>

Tel: 0151 705 3220

Fax: 0151 705 3241

UKAS ACCREDITED FOR TESTS PUBLISHED ON SCHEDULE OF ACCREDITATION. REFERENCE 9362.

Please see UKAS website for further details on scope of practice. Details also listed within section 4 of this handbook.

CONTENTS		PAGE
1	INTRODUCTION	4
	1.1 Introduction to the laboratory	4
	1.2 Quality policy	4
	1.3 Using this handbook	5
	1.4 COVID-19 pandemic	5
2	LABORATORY INFORMATION	6
	2.1 Opening times	6
	2.2 Turnaround times	6
	2.3 Contact details	6
3	ADVICE FOR USERS	7
	3.1 Specimen containers	7
	3.2 Request forms	7
	3.3 Patient identifiers / Specimen acceptance	7
	3.4 Packaging and Transport	8
	3.5 Urgent requests / High risk specimens	8
	3.6 Specimen retention times	9
	3.7 Protection of personal information policy	9
	3.8 Complaints procedure	9
	3.9 Service agreement	9
4	TESTS OFFERED AND KEY FACTORS AFFECTING TESTS	10
	4.1 Blood Parasite diagnosis	10
	4.1.1 Malaria	10
	4.1.2 African Trypanosomes	11
	4.1.3 Filariasis	11
	4.2 Faecal Parasite diagnosis	12
	4.2.1 Faecal microscopy (including hot stool)	12
	4.2.2 Culture for Strongyloides / Hookworm	12
	4.2.3 Faecal staining methods	13
	4.2.4 Enterobius microscopy	13
	4.2.5 Faecal PCR	14
	4.3 Urine techniques	14

4.4	Leishmania diagnosis	15
4.4.1	Microscopy	15
4.4.2	PCR	15
4.5	Antibody detection	16
4.5.1	Amoebiasis	16
4.5.2	Malaria	17
4.5.3	Leishmaniasis	17
4.5.4	African Trypanosomiasis	18
4.5.5	South American T.cruzi	18
4.5.6	Schistosomiasis	19
4.5.7	Strongyloidiasis	19
4.5.8	Hydatid	20
4.5.9	Filariasis	21
4.5.10	Fasciola	21
4.6	Other samples	22
4.6.1	Semen	22
4.6.2	Sputum	22
4.6.3	Histology	22
4.6.4	Insects	22
4.6.5	Whole worms / proglottids	22
4.6.6	Amoebiasis	23
4.6.7	Cyst fluid for Hydatid disease (<i>Echinococcus</i>)	23
5	CHARGES	23
6	RESULTS AND REPORTS	23
6.1	Written reports	23
6.2	Telephoned / faxed reports	24
6.3	Interpretation and advice (including out of hours)	24
	Appendix 1	25

1 INTRODUCTION

1.1 Introduction to the laboratory

The Clinical Diagnostic Parasitology laboratory (CDPL) is a UKAS accredited laboratory working to ISO15189:2012 standard, accredited for all tests placed on scope of practice, see section four of this handbook for details of tests accredited by UKAS. The CDPL is located in the Department of Clinical Sciences at The Liverpool School of Tropical Medicine. The laboratory offers a referral service for the identification of a wide range of human parasites from clinical specimens. The laboratory participates in the external NEQAS quality assurance schemes for blood parasitology, faecal parasitology, malaria rapid diagnostic tests (RDT's) and parasite serology.

The laboratory accepts samples from NHS and PHE laboratories, private hospitals, and private laboratories.

The laboratory cannot accept samples directly from patients; samples must be referred to the laboratory by a health professional.

1.2 Quality policy

Our Quality Policy presented below aims to offer a first-class nationwide service for the laboratory diagnosis of parasitic infections.

The Clinical Diagnostic Parasitology Laboratory is committed to providing a high-quality diagnostic service responsive to the needs and requirements of its users and will:

- Provide a high-quality diagnostic parasitology service encompassing; blood and faecal parasitology, parasite serology and other diagnostic parasitology tests according to individual patient needs as appropriate.
- Operate a quality management system to integrate the procedures, processes and resources of the services provided.
- Set quality objectives and plans in order to implement this quality policy.
- Ensure that all personnel are familiar with the quality manual and all policies and procedures relevant to ensure user satisfaction.
- Commit to the health, safety and welfare of all staff.
- Treat visitors to the unit with respect and provide for their safety.
- Maintain good professional practice and conduct including compliance with relevant environmental legislation.

The Clinical Diagnostic Parasitology Laboratory will comply with standard ISO15189:2012 and is committed to:

Page 5 of 25

- Staff training and development to provide an appropriate and effective service to users.
- Procurement and maintenance of equipment and resources necessary for provision of the service.
- Appropriate handling of specimens to ensure correct performance of laboratory examinations.
- Examination procedures that will ensure the high quality of all tests offered.
- Reporting results in a timely, confidential, accurate and clinically useful manner.
- Regular assessment of user satisfaction, in addition to internal audit and external quality assessment, in order to produce continual quality improvement.

Signed on behalf of the Clinical Diagnostic Parasitology Laboratory, LSTM,



19.05.20

Prof Bertie Squire, Dean of Clinical Sciences and International Public Health
Clinical Director



19.05.20

Mrs Jayne Jones BSC, MSC, MIBMS
Laboratory Manager

1.3 Using this handbook

This handbook aims to provide help and advice for users of our service. Information contained in this handbook should help the user understand the requirements needed for the diagnostic parasitology services offered by the laboratory. Also included is information regarding interpretation of results. This handbook should assist with pre-examination and post-examination information needed by the user. If users require any additional information that is not provided in this handbook, then they should contact the laboratory for advice.

1.4 COVID-19 pandemic

The CDPL continues to offer a full service during the COVID-19 pandemic. Serology turnaround times have been extended slightly to allow for fewer staff being onsite at any one time for social distancing purposes. Urgent testing can still be requested.

2 LABORATORY INFORMATION

2.1 Opening times

Please note that the Laboratory is open between the hours of 8am to 5pm Monday to Friday. Specimens received after 4pm may not be processed until the following day.

The laboratory is closed at weekends and on Bank Holidays.

The laboratory does not provide an on-call / out of hours' service.

The laboratory is a department of LSTM which is closed over the Christmas and New Year period therefore a limited service is offered. Christmas closures are sent to users informing dates of closure.

2.2 Turnaround times

TEST REQUEST	MAXIMUM TURNAROUND TIME
Malaria / other blood parasite microscopy	Same day
Blood filtrations	Same day
Faecal parasite microscopy	2 working days
Faecal parasite staining	3 working days
Faecal parasite PCR	7 working days
Strongyloides stool culture	7 working days
Urine microscopy / filtration	2 working days
Parasite antibody serology tests	10 working days
Leishmania PCR	7 working days (if speciation required up to 10 working days)
Leishmania microscopy	3 working days

Turnaround times for some additional tests are variable – please contact the laboratory for advice.

2.3 Contact details

Please see the information on page one of this handbook for all contact details.

3 ADVICE FOR USERS

3.1 Specimen containers

The majority of procedures used in this laboratory require standard specimen containers such as plastic stool pots and plastic serum tubes.

Leishmania PCR is the only test where use of buffer and Eppendorf tubes supplied by our laboratory are preferred. Information is available under the “Leishmania” section of this handbook (page 13).

3.2 Request forms

It is essential that a request form accompanies all samples. The current version of our request form can be found on the internet, address on page one of this handbook.

All available patient identifiers must be completed along with hospital reference numbers. Clinical details and travel history are helpful regarding diagnosis and interpretation and should be completed where possible. The requesting laboratory must include contact details on the form for return of results. Test requested and type of sample must be completed along with the priority status and high-risk information.

If users send their request form it must contain all information required such as patient identifiers, sending location, clinical details, sample sent, and test requested.

3.3 Patient identifiers / Specimen acceptance

In order to comply with specimen acceptance policy, samples sent to the laboratory for diagnosis plus accompanying forms must be labelled with a minimum of 3 out of the 6 sample identification criteria:

1. Family name (or reference number)
2. First name(s)
3. Date of Birth
4. Referring laboratory reference number
5. Hospital number
6. Specimen date

Unfortunately, specimens not stating at least 3 of these criteria will be rejected.

3.4 Packaging and Transport

Specimens must be packaged in compliance with UK postal regulations for Diagnostic Biological samples – UN3373

Specimens should be in an appropriate container surrounded by absorbent material and then placed in a leak proof sample bag. The request form should be placed in a separate pocket of that bag; all should be placed inside a plastic container which is then placed into an appropriate cardboard box. The cardboard box should be labelled with recipient's full address along with senders contact details. The box should be clearly marked as UN3373 – Biological substance category B, pathological specimen fragile with care.

Specimens not packed in this way may leak in transit and be unsuitable for testing.

Samples should be sent by first class mail, or courier to:

The Clinical Diagnostic Parasitology Laboratory
Liverpool School of Tropical Medicine
Pembroke Place
Liverpool
L3 5QA

OR

If in the Hays DX scheme:
Liverpool School of Tropical Medicine
Diagnostic Laboratory
DX6966301
Liverpool 92L

The sender is responsible for ensuring health and safety of any courier or taxi service that is used to transport specimens to the laboratory.

3.5 Urgent requests / High Risk specimens

Please ensure that the laboratory is informed if samples are High Risk and / or in need of urgent testing. The laboratory cannot process samples from patients with TB or VHF. The laboratory does not knowingly deal with specimens from patients with Category 3 infections such as viral haemorrhagic fevers.

Urgent testing can be performed if required. If this results in a test being performed in addition to its routine timescale, for example antibody testing, then an additional cost will be incurred. Requests must be marked urgent and ideally the laboratory telephoned in advance.

3.6 Specimen retention times

If there is need of additional testing of a sample already sent to the Diagnostic Parasitology laboratory, please note the following sample retention times:

- EDTA blood - 1 day
- Faecal samples - 2 days
- Serum samples - 1 year
- Negative blood films – 3 months
- Positive blood films – 1 year

3.7 Protection of personal information policy

The Clinical diagnostic parasitology laboratory has a storage of information policy that ensures patient data and information, either electronically or as a hard copy, is kept confidential and secure at all times. All levels of staff adhere to this policy. There is a local privacy notice for all users, available on the CDPL website.

3.8 Complaints procedure

Any complaints received by the Clinical Diagnostic Parasitology Laboratory are dealt with as a matter of urgency. The laboratory has a written policy on dealing with complaints. Please contact the laboratory manager if a complaint arises.

3.9 Service agreement

The Clinical Diagnostic Parasitology Laboratory (CDPL) at the Liverpool School of Tropical Medicine considers every request received for a test carried out within our department as a service agreement between the requestor and the CDPL. Once a specimen is accepted for testing a service agreement is established and the specimen processed according to appropriate SOP's. The service agreement will be reviewed upon receipt of sample and during issue of results. The request, specimen received, suitability of test carried out, recommendations for further tests, interpretation of result and the final report all constitute part of that service agreement review. Tests offered in the CDPL have been validated and

evaluated for use in the unit. The CDPL management are confident that the tests offered are appropriate to the service provided. Staff employed by the CDPL are in roles specific to their qualifications and HCPC registration and regular staff competency testing is carried out.

4 TESTS OFFERED AND KEY FACTORS AFFECTING TESTS

If a sample is received for a test the laboratory does not offer the user will be contacted and if requested the sample forwarded to the relevant laboratory.

If the sample has been home collected by the patient the date and time of specimen still needs to be recorded. Please see information for each individual test regarding sample age and storage procedures.

Specimens will be rejected if they do not meet any sample requirements for the test requested.

4.1 Blood Parasite diagnosis - UKAS accredited test

4.1.1 Malaria

Samples are screened using thick blood films, thin films are examined on positive samples. If *P.falciparum* is found, parasitaemia is performed. Rapid diagnostic tests (RDT's) are available.

Sample requirements:

- Thick and thin films (unstained but thin film methanol fixed) made from fresh blood sample.
- Thick and thin films made and stained from fresh blood sample.
- Original EDTA sample.

Key factors affecting tests: malaria parasites collected into anticoagulants such as EDTA deteriorate and morphological changes occur within a few hours.

We would be grateful if the malaria reference laboratory form could be completed, along with our report form, and enclosed with the sample. Sample date and patient travel history should be stated. Malaria reference laboratory form is available on the internet; address can be found on page one of this document.

4.1.2 African Trypanosomes

Thick blood films and concentration technique if films are negative. Blood samples for concentration should be examined on the day of collection.

Sample requirements:

- unstained thick/thin films
- EDTA blood sample

Key factors affecting tests: Trypanosomes may not survive for more than a few hours in EDTA tube therefore posted samples could show false negative results. **PHONE LAB FOR ADVICE BEFORE SENDING SPECIMEN.**

If a CSF sample is to be examined for trypanosomes it should ideally be examined within 20 minutes of L.P.

4.1.3 Filariasis

A definitive diagnosis of filariasis is usually made by the demonstration of microfilariae in the peripheral blood. The exception to this is the diagnosis of *Onchocerca volvulus* which is diagnosed by demonstration of microfilariae in skin snips.

Blood filariasis:

A wet preparation is examined and samples are filtered using a Nuclepore membrane. If *W.bancrofti* is suspected blood collection time should be between 22.00 and 02.00 hours, for *Loa loa* diurnal blood sample (noon) is preferred.

Sample requirements: EDTA blood. Sample size is not critical but ideally between 5ml and 10ml should be sent, the larger the sample filtered the greater chance of demonstrating the microfilariae.

Key factors affecting tests: sheathed microfilariae may ex-sheath if blood sample is not examined within 2 to 3 days of collection, all microfilariae may last up to 72 hours in EDTA blood before disintegration.

Onchocerca volvulus:

Please contact the laboratory before sending skin snips for microfilariae diagnosis, contact details on page one of this handbook.

4.2 Faecal Parasite diagnosis

4.2.1 Faecal microscopy (including hot stool) - UKAS accredited test

A concentration technique is routinely performed on all faecal samples for the presence of cysts, ova and larvae. Direct saline smears are examined on samples that are 24 hours old or less where trophozoite stages of protozoan parasites are suspected.

Sample requirement: Minimum of ¼ specimen pot of faecal sample. Ova, cysts and parasites may be passed intermittently therefore three samples may be required to be examined. Ova, cysts and parasites will diminish over time therefore it is more ideal for the sample to be less than 2 days old upon receipt at LSTM, however older stools will not be rejected.

Key factors affecting test: trophozoites may only survive for up to 24 hours in voided faecal samples

If *Entamoeba histolytica* trophozoites are suspected a “hot” stool must be examined.

Hot stool sample requirement: specimen must reach the laboratory within 30 minutes of sample being produced and must be marked as hot stool, this will then be examined with priority. Methanol fixed faecal smears can be made at requesting site if the sample can not arrive at CDPL within 30 minutes, these smears can then be sent to CDPL for staining and examination.

4.2.2 Culture for Strongyloides / Hookworm - UKAS accredited test

Faeces are cultured for filariform larvae using the charcoal technique.

Sample requirements: At least 50g (half pot) of faeces is required for this test. Ova, cysts and parasites may be passed intermittently therefore three samples may be required to be examined.

Key factors affecting tests: samples should **NOT** be stored in a refrigerator/cold room following collection as this may inhibit subsequent larval growth.

4.2.3 Faecal staining methods - UKAS accredited test

Cryptosporidia

Slides from concentrated faecal deposit are stained using the Z-N technique for identification of oocysts.

Sample requirement: minimum of ¼ pot of faecal sample. Ova, cysts and parasites may be passed intermittently therefore three samples may be required to be examined.

***Dientamoeba fragilis* – This subsection NOT UKAS accredited test**

Romanowsky stained direct smears are examined for the identification of *D.fragilis* trophozoites.

Sample requirements: please send a fresh (dated) unfixed sample of faeces (minimum of ¼ pot) plus a **thin** faecal smear -made without saline and fixed once dry for 1 minute in methanol/IMS

Key factors affecting test: ideally sample should not be more than 1 day old when sent for testing as trophozoites degenerate.

4.2.4 *Enterobius vermicularis* microscopy

The rectal swab method is used in preference to sellotape slides, however both means of sample collection will be examined.

Sample requirements: a cotton bud dampened with physiological saline is wiped around the peri-anal area, and placed in a small bottle of physiological saline. The bottle is sent to the laboratory for examination. OR adhesive tape is applied to the area, attached sticky slide down to a microscope slide and sent to the laboratory for examination.

Key factors affecting test: ideally the swab / sellotape sample should be taken in the morning before washing the peri-anal area.

4.2.5 Faecal PCR

We have recently introduced a qPCR technique for intestinal parasites. The parasites included in the PCR panel are; *Giardia intestinalis*, *Cryptosporidium* species, *Entamoeba histolytica*, *Entamoeba dispar*, *Strongyloides stercoralis*, *Necator americanus*, *Ancylostoma duodenale*, *Schistosoma* species, *Ascaris lumbricoides* and *Trichuris trichiura*.

From our intensive evaluations we have concluded that one stool sample received and tested using stool concentration, faecal PCR and helminth culture (if required), will yield the same results as the recommended three stool sample request for stool concentration and helminth culture (if required). The introduction of the faecal PCR will therefore eliminate the need for requesting three stool samples from patients for parasitology testing and move to a one stool sample request. This will be of benefit to the patient plus enhance the sensitivity and confidence in results from a one-off stool sample test for users of our service.

If you have any clinical concerns about this new approach, please contact one of the consultant physicians via on call mobile telephone 07909910899. This telephone is in operation Monday to Friday 9am to 5pm.

Sample requirements: Minimum of ¼ specimen pot of faecal sample.

Key factors affecting test: Addition of formalin to the sample will hinder DNA extraction and qPCR amplification.

Appendix 1 on page 25 of this document shows the sensitivity and specificity values for this test.

4.3 Urine techniques - UKAS accredited test

Schistosoma haematobium

Urine is tested by dipstick for the presence of blood /Hb, and RBC's. Urine is routinely filtered for Schistosomiasis ova using polycarbonate filters.

Sample requirement: ideally total 10am – 2pm urine collection or terminal midday sample

Key factors affecting test. Random urine samples may give false negative results

4.4 Leishmania diagnosis

4.4.1 Microscopy- UKAS accredited test

4.4.2 PCR

Sample requirements: See downloadable sampling notes on website, address can be found on page one of this document.

Cutaneous Leishmaniasis

- Unstained, fixed aspirate or biopsy impression smears for microscopy.
- Giemsa-stained histology slides.
- Biopsy material in PCR (ATL) buffer (available from laboratory on request), OR in dry, sterile container OR in 10% ethanol. If histological wax block of tissue is the only material available please cut 10µm thick wax sections and float onto microscope slides (3/slide, two or 3 slides/sample), send these unstained for DNA extraction. A biopsy should ideally be around the size of a grain of rice.

Visceral leishmaniasis

- THIN marrow smears (please fix for 1 minute in methanol before sending.)
- Marrow /blood in EDTA for PCR -minimum of 500µl

Key factors affecting test: Use of aspirate, wax block or samples smaller than the size of a grain of rice MAY cause false negative or insufficient DNA results. The use of Iodine during sampling will inhibit the PCR amplification and may cause false negative results. The use of Lithium heparin instead of EDTA will inhibit the PCR amplification and may cause false negative results. Please list travel history with all Leishmania PCR requests.

4.5 Antibody detection

Interpretations of all results are subject to clinical and travel information provided.

Antibody detection may not be useful as a test of cure.

A positive result may reflect previous or current infection. We can examine samples for parasites following a positive antibody test result if required.

Different serology tests have differing cut off points therefore more information regarding tests and interpretation is available from the laboratory staff.

Sample requirement: 0.5ml serum OR 10ml of clotted blood (minimum 0.1ml serum or 5ml clotted sample)

Key factors affecting tests. Serum samples ideally should not be sent “on the clot”. Please use serum gel or similar tubes to avoid haemolysis of serum if delayed in transit as there is a possibility that the test result may be affected/invalid due to storage temperature. Plasma can be tested, although due to the fact that there are more proteins present in plasma samples than in serum samples the risk of an incorrect result is increased.

Please ensure geographical location / travel history has been stated on the form if known, this is particularly important for Trypanosomiasis testing.

4.5.1 Amoebiasis (Amoebic Liver abscess) - UKAS accredited test

IFA used as initial screen, urgent or positive IFA samples will have a secondary test performed. Up to 95% of Amoebic liver abscess (ALA) patients are sero-positive by two weeks' post infection. Lower level positive results may be obtained if tested earlier than two weeks' post infection. In these cases, a low level positive result may be significant and a repeat test should be performed at 14 days' post infection. Titre will decrease in 1 to 2 months after successful treatment but may remain low level positive for an extended time.

Entamoeba histolytica antigen used. IgG antibodies detected.

	SENSITIVITY %	SPECIFICITY %
IFA AMOEBIASIS	93.8	96
SECONDARY TEST (in house verification)	100	91.6

4.5.2 Malaria

IFA used.

If current infection is suspected blood films must be examined. Antibody testing may be useful for retrospective diagnosis.

Antibodies are detected about 7 to 15 days from initial infection.

In a non-immune traveller treated for a single infection Ab levels should fall within 3-6 months of successful treatment.

Plasmodium falciparum and *Plasmodium vivax* antigens available, however species are known to cross react. *Plasmodium falciparum* antigen used unless otherwise requested. IgG antibodies detected.

4.5.3 Leishmaniasis

IFA used plus commercial Leishmania dipstick in non-cutaneous queries.

Ab levels detected around 1 month post exposure.

The IFA should seroconvert to negative within 6-9 months of successful treatment.

Antibody detection is useful in cases of suspected Visceral Leishmaniasis who do not have HIV infection. In suspected Visceral Leishmaniasis patients who are HIV infected sera is often non-reactive.

In Cutaneous Leishmaniasis, serology is of little value.

Cross-reactions may occur with trypanosomiasis and autoimmune conditions.

	SENSITIVITY %	SPECIFICITY %
LEISHMANIA IFA	94	94
LEISHMANIA DIPSTICK (in house verification)	75	100

IFA uses *Leishmania donovani* antigen. IgG antibodies detected. Secondary dipstick detects rK39 antigen.

4.5.4 African Trypanosomiasis

IFA used for *T.b.rhodesiense* queries. IgG antibodies detected.

CATT used for *T.b.gambiense* queries:

In-house verification SENSITIVITY 100%

In-house verification SPECIFICITY 98%

Travel history and clinical details are required to determine which test is performed.

Cross reactions may occur between trypanosome & leishmania serology.

CSF samples can also be tested for African trypanosomiasis antibodies.

4.5.5 South American T.cruzi - UKAS accredited test

Serology is useful, Ab's are detected around 1 month post- infection and remain positive for life.

Commercial T.cruzi test used.

	SENSITIVITY %	SPECIFICITY %
in house verification	100	100

Positive results are always repeated for quality purposes, firstly at LSTM, then LSTM clinician would recommend user to obtain a repeat test at another site.

4.5.6 Schistosomiasis - UKAS accredited test

Commercial ELISA used. Test utilises *Schistosoma mansoni* antigen, however this does cross react with other *Schistosoma* species. IgG antibodies are detected.

	SENSITIVITY %	SPECIFICITY %
ELISA (in house verification)	100	60

Antibody levels may not be detected until 3 months' post exposure. In patients with a negative ELISA test up to 3 months but significant fresh water exposure we would recommend further testing from 4 months after the last exposure.

Antibody levels may be positive for 18 months or longer post successful treatment.

4.5.7 Strongyloidiasis - UKAS accredited test

Commercial ELISA used. IgG antibodies are detected.

Cross-reactions are known to occur with patients with filarial infections and with heavy Hookworm infections.

HIV status has no effect on Strongyloides antibody test

Antibody levels normally detected 2 to 3 months post exposure, however, there are occasions when no antibody response will be detected.

Antibody levels revert to negative approximately 6 months to 1 year post successful treatment.

	SENSITIVITY %	SPECIFICITY %
ELISA (in house verification)	95	95

4.5.8 Hydatid (*Echinococcus granulosus*) - UKAS accredited test

Samples are screened with commercial ELISA. IHA is performed on ELISA positive samples.

Amongst proven cases of hydatid disease, 92% show a positive ELISA test. Sensitivity depends on cyst site: Liver 96%, pulmonary 76%, skeletal 60%, other sites vary. Brain hydatid rarely shows positive serology. Serological cross-reactions, giving rise to false positives, can occur with other parasitic infections, particularly larval cestodes and filarial worms and with some neoplasms. Less than 3% of non-infected controls are positive.

False negatives may occur (about 8%) and are more common in patients with extra-hepatic cysts. False negatives can be due to calcified cysts. Patients with cysts occurring in the brain are usually serologically negative.

Antibody levels are detected at variable timescales post infection due to nature of disease.

Antibody levels may remain positive for life post successful treatment / surgery.

IHA is performed on all samples with positive hydatid ELISA serology

	SENSITIVITY %	SPECIFICITY %
ELISA (in house verification)	100	97.6
IHA (manufacturers analysis)	93	94.9

The sensitivity of the ELISA is estimated to be 97%, the cases with negative serology but subsequently proven hydatid disease have mainly been those patients with extra-hepatic lesions.

Samples found equivocal by ELISA are re-tested by IHA. We consider results significant if both tests are positive. If sample shows equivocal results, we would suggest repeating serology in 1 months' time.

ELISA utilises *Echinococcus* species antigen and detects IgG antibodies.

IHA uses *Echinococcus granulosus* antigen.

4.5.9 Filariasis - UKAS accredited test

Commercial ELISA used. This utilises *Acanthocheilonema viteae* antigen, that will cross react with human filariasis species, and detects IgG antibodies.

Antibody levels detected around 2 to 3 months’ post exposure.

Antibody levels negative from 1 year or longer post successful treatment.

	SENSITIVITY %	SPECIFICITY %
ELISA (in house verification)	100	90

4.5.10 Fasciola - UKAS accredited test

Commercial ELISA used. This utilises *Fasciola* species antigen and detects IgG antibodies.

Antibody levels detected around 2 to 3 months’ post exposure.

Antibody levels should revert to negative around 6 months’ post successful treatment.

	SENSITIVITY %	SPECIFICITY %
ELISA (In house verification)	100	77

4.6 Other samples

4.6.1 Semen - UKAS accredited test

Seminal fluid microscopy is performed for schistosome ova.

Specimen requirement: Age and volume of sample is not critical.

4.6.2 Sputum

Sputum microscopy is performed for parasitic ova.

Specimen requirement: Age and volume of sample is not critical. The laboratory does not process samples when a diagnosis of TB cannot be excluded.

4.6.3 Histology

Histology sections should be sent stained for parasite microscopy. Ideally H&E stained sections plus Giemsa stained section if querying Leishmaniasis or a PAS stained section if querying Amoebae.

4.6.4 Insects

Specimens query insects or ectoparasites must be sent in a clean dry container – not attached to sellotape.

4.6.5 Whole worms / proglottids

Identification of whole worms/proglottids

Sample requirements: Whole worms such as Ascaris may be stored in 10% formol saline. Suspected Taenia species proglottids are best stored in **physiological saline** and sent to the laboratory as soon as possible. The use of 10% formol saline should be avoided unless the sample cannot be sent to the reference laboratory within 3 days.

Key factors affecting tests: Identification may be difficult if formalin-fixed proglottids are sent for identification, however proglottids may disintegrate if in saline for 3 days or more.

4.6.6 Amoebiasis – examination of cyst aspirate / pus samples **UKAS accredited test**

Specimen requirement: Requests for *Entamoeba histolytica* in samples of pus from liver abscesses should either be examined within one hour of collection or fresh THIN smears of pus should be made (minimum of 3), fixed when dry (1 minute in methanol) and sent for staining together with pus sample.

Key factors affecting test: About 30 minutes after aspiration of cyst pus/fluid *E.histolytica* trophozoites become indistinguishable from macrophages.

4.6.7 Cyst fluid for Hydatid disease (*Echinococcus* species) **UKAS accredited test**

Aspiration of cyst should only be considered after taking expert advice.

Specimen requirement: Cyst fluid should ideally be examined within 2 days of sample being taken. No minimum sample requirement.

5 CHARGES

A charge is made for all laboratory services, for current prices please telephone laboratory staff on 0151 705 3220

6 RESULTS AND REPORTS

6.1 Written reports

Reports are processed and issued as soon as results are available and have been authorised. Copies can be issued if required please contact the laboratory if this is needed.

6.2 Telephoned / faxed reports

Positive malaria film results are telephoned as soon as a diagnosis is available. If the report is to be telephoned to a specific individual, then please add their details to the request form.

Other results may be telephoned if the laboratory manager considers it appropriate.

Other requests for telephoned results are reported when a result is available. Please state on request form if telephoned results are required.

Once a result has been telephoned, a hard copy is also sent, and it will be indicated on the form that the result has already been telephoned.

Results may be faxed if requested. The user must have access to a secure fax, patient names are not included on faxed transmissions but all other identifiers remain.

6.3 Interpretation and advice

For technical advice regarding samples/tests and for interpretation of results please contact the diagnostic laboratory manager 0151 705 3220/3290

For clinical advice on treatment of parasitic diseases please contact one of the consultant physicians via on call mobile telephone 07909910899. This telephone is in operation Monday to Friday 9am to 5pm. Out of hours this number will transfer to Infectious diseases unit at the Royal Liverpool University Hospital. If any issues occur regarding transfer of call out of hours, please telephone the RLUH switchboard and ask for Infectious diseases 0151 7062000.

Appendix 1 – sensitivity and specificity of faecal PCR

Parasite	Sensitivity %	Specificity %
<i>Cryptosporidium</i> species	100	100
<i>Entamoeba histolytica</i>	65	100
<i>Giardia intestinalis</i>	83	91
<i>Ascaris</i>	100	100
<i>Trichuris</i>	93	97
<i>Schistosoma</i> species	NA	99
<i>Strongyloides</i>	100	98
<i>Hookworm</i>	88	98